

Fundamental Study to New Evaluation Method Based on Physical and Psychological Load in Care

Hiroaki Inoue

Tokyo University of Science, Suwa
Research course of Engineering/Management
Chino-city, Japan

Hiroyuki Nara

Hokkaido University
Graduate School of Information Science and
Technology
Sapporo-city, Japan

Fumikazu Miwakeichi

The Institute of Statistical Mathematics
Spatial and Time Series Modeling Group
Tachikawa-city, Japan

Senichiro Kikuchi, and Satoshi Kato

Jichi Medical University
Department of Psychiatry
Shimotsuke-city, Japan

Shunji Shimizu, Hiroataka Ishihara and Yuuki
Nakata

Tokyo University of Science, Suwa
Department of Computer Engineering
Chino-city, Japan

Takeshi Tsuruga

Hokkaido Institute of Technology
Department of Clinical and Rehabilitation Engineering
Sapporo-city, Japan

Nobuhide Hirai

Tokyo Medical and Dental University
Health Service Center
Bunkyo-ku, Japan

Eiju Watanabe

Jichi Medical University
Department of Neurosurgery
Shimotsuke-city, Japan

Abstract—In Japan and developed countries, it has become aged society, and wide variety welfare device or system have been developed. But these evaluation methods of welfare device or system are limited only stability, intensity and partial operability. Because of, it is not clear to determine the standard to evaluation for welfare device or system of usefulness. Therefore, we will attempt to establish the standard for evaluation about usefulness for objectively and quantitatively for including non-verbal cognition. We examine the relationship between human movements and brain activity, and consider the evaluation method of welfare devices and systems to measure the load and fatigue which were felt by human. In this paper, we measure the load for sitting and standing movement using NIRS. We tried to make sure for the possibility of the quantitatively estimation for physical or psychological load or fatigue by measuring of brain activity using NIRS(Near Infra Red Spectroscopy). As results, when subjects perform the movement task, the statistical significant difference was shown in the specific part of the brain region.

Keywords—component; Evaluation; Movement; Exercise; NIRS; Care; Welfare Technology; Useful welfare device evaluation; Evaluation method

I. INTRODUCTION

As it has been known widely, aging population in Japan and world-wide countries has been increasing. Thereby the

number of care worker has been increasing. Care is very hard work. Welfare devices and systems reducing a burden of the care work are required. In this background, welfare systems and device are rapidly developing, and various devices are manufactured based on the increased popularity of welfare device and system. Also, the market of welfare devices and systems is expanding. However, the evaluation method is limited respectively to stability, strength and a part of operability for individual system or device. It means that evaluation methodology for usefulness of them was not established. Therefore, we will attempt to establish a standard to evaluate the usefulness for objectively and quantitatively on the basis of cognition such as physical load, reduction of fatigue and postural stability. Especially, in considering universality, it is necessary to measure human movement in daily life. Movement was not measured by using particular device, but routinely-performed movement in daily life.

In our previous study, we examined the possibility of evaluation by measuring physical load due to activities of daily living with using 3D Motion Analysis System and EMG. Also, we looked into the possibility of quantitative evaluation of tiredness and load on the basis of brain activity using NIRS [1]. We consider that physical and psychological load are linked to cognition including non-verbal cognition. In this paper, our purpose is to obtain an index of the measure of load and fatigue which are felt by human. We discussed

experiments which explore the relationship between exercise and human brain activity. In addition, we measured the brain activity of subjects who performed exercise task with imposing additional load. Thus, we tried to measure the load and fatigue felt by human.




Fig. 1. Experimental View by using NIRS

II. EXPERIMENTAL METHOD

A. Evaluation by using NIRS

We measured the subject's movement and muscle potential in our previous research [1]. In this paper, we would like to discuss about brain activity when subjects performed the movement tasks which are standing and sitting (Figure 1).

Subjects were six males aged twenty. They were asked to read and sign an informed consent regarding the experiment. Measurement apparatus was NIRS (SHIMADZU CO. Ltd products-FOIRE3000 [4]). Measurement region was at right and left prefrontal cortex.

1) Measuring brain activity during transfer with standing position (task1)

At this measurement, the subjects used welfare device to perform transferring in a standing position. In this measurement, subject sat on seating face of welfare device appeared on the top of chair after raising hip until kneeling position. Also, subject performed inverse transferring from seating face to chair. Time design was rest (5 seconds), task (10 seconds), and rest (5 seconds). This time design was repeated 30 times. Rest time is to stabilize the brain activity. In the measurement NIRS,

2) Measuring brain activity during transfer with half-crouching position (task2)

At this measurement, the subjects used welfare device to perform transferring in a half-crouch position. In this measurement, the subjects sat on seating face of welfare device appeared on the top of chair after raising hip until kneeling position. Also, the subject performed inverse transfer from seating face to chair. Time design was rest (5 seconds), task (10 seconds) and rest (5 seconds). This time design was repeated 30 times.

In experiments of task1 and task2, the operation of welfare device was performed by an operator other than subject. Before this measuring, subjects adjusted to transferring by use of welfare device.

B. Measuring the brain activity when subjects perform movement task with additional load.

In this experiment, we have performed measuring brain activity in the case where no load is applied to subject. In previous experiment, load of subjects were derived from only standing and sitting movement. In this experiment, we added the weight to subjects as additional load. As load, there are some cases which are no load 5kg and 10kg. Subjects wear a backpack containing the weight. And Subjects performed standing and sitting movement. Task design was same with previous experiments.

III. EXPERIMENTAL RESULTS

A. Evaluation by using NIRS

As the common result of all subjects, oxy-Hb tended to increase during task and to decrease in resting state. Therefore, it was thought that change of hemoglobin density due to task was measured. Fig.3 and Fig.4 show trend of the channel in which significant different was shown. Analysis was performed via one-sample t-test [5,6,7,8,9] by a method similar to previous researches [5,6,7,8,9]. In this analysis, it was necessary to remove other than change of blood flow due to fatigue. So, our method was mainly focused on resting state to compare with the 1st trial and other trials of brain activity.

In task1, 1 and 2, each of sample data for analysis was 4 seconds after the task (Fig. 2). In the t-test of the same task, we performed t-test with first time trial and other trial which was from second times to thirty times, and examined relationship the number of trials and significant differences.

In task 1, significant different could be found from the about 10th trials. Fig. 5 show the region confirmed significant difference. In task 2, significant different could be found from the about 10th trials too. Fig. 6 shows region confirmed significant difference.

At first, we performed t-test using 4 seconds during first trial and 4 seconds during other trials, which were from second to fifteenth in same position.




Fig. 2. T-test of sample data in task1 and 2




Fig. 3. Measuring result of task1




Fig. 4. Measuring result of task2




Fig. 5. Significant difference of task1




Fig. 6. Significant difference of task2

B. Results of Brain activity measurements when the load was applied to the subjects

Analysis method was one-sample t-test of brain activity data as with above analysis. Fig.7 shows the result of one sample t-test between first trial of rest data and another trial of rest data. This analysis method was same with Fig.2. Fig.8 and Fig.9 shows the result of one sample t-test with brain activity data of different movement in the same number of trials. Red circle is the brain region that has seen a statistically significant difference in 5 of 6 subjects. Yellow circle show the brain region that has seen a statistically significant difference in 4 subjects. Green circle show the brain region that has seen a statistically significant difference in 3 subjects.

In the both t-test, there were significant difference on the prefrontal cortex. These results were similar to above experiments. However, significant differences were found randomly regardless of the number of trials. As the cause of these results, there is a possibility that t-test could not remove the changes in scalp of blood flow.




Fig. 7. Result of one sample t-test between first trial and another trials when subjects had no additional load




Fig. 8. Result of one sample t-test between first trial and another trials when subjects had 5kg load.




Fig. 9. Result of one sample t-test between first trial and another trials when subjects had 10kg load

IV. DISCUSSION

A. Evaluation by using NIRS

In this experiment, we tried to measure quantitatively the physical and psychological strain on the basis of brain activity. Also, we think that brain activity disclose human cognitive including non-verbal. As a result, it was shown that there were differences at brain activity due to number of trials and postural.

In this time, analysis was performed via one-sample t-test using sample of brain activity in resting state during task or after task. Hence, analysis method was to remove disturbance such as body motion and angular variation of neck to the extent possible although there was the possibility to measure skin blood flow. Therefore, it was thought that strain due to tasks was quantitatively measured by being recognized significant differences. Also, in previous research, it was reported to decrease in activity in the brain around #10, 11 [10], as the result of measuring brain activity during Advanced Trial Making Test using PET [11]. Therefore, this result came out in support of previous research in no small part.

B. Results of Brain activity measurements when the load was applied to the subjects

As a step to make the evaluation method, we performed additional experiment, which is imposed to subjects the load other than using the welfare devices. Significant difference was observed in the brain region similar to previous experiments. And, each time the load increases, brain regions found statistically significant differences became widespread. And, the frequency of showing the statistical significant difference became the higher. We think that there are the possibility of happen this results by the additional load.

Of course, it is necessary to increase number of subject at the present stage. In addition, there are problems associated with experiment, number of subject, method and measured region. However, in terms of being recognized significant differences at brain activity due to movement, it was thought to show useful result in evaluating quantitatively daily movements.

V. CONCLUSION

In this paper, we tried to measure physical and psychological load with measuring brain activity. And there were significant differences due to number of trials. In this experiment, analysis method was to remove disturbance such as body motion and angular variation of neck to the extent possible by using the measurement result in resting state as sample. Therefore, it was thought to show the useful result in evaluating quantitatively load due to movement task by being recognized difference in brain activity caused by number of trials.

Main purpose in this study is to evaluate physical load and fatigue quantitatively. So, we tried to evaluate change of muscle load due to difference of motion by simultaneous measuring with 3D motion analysis System and EMG quantitatively.

However, evaluation of psychological load is necessary, too. In terms of using welfare device, prolonged use must be taken into account. In this case, it is important to consider not only physical load but also psychological load due to prolonged use from standpoint of developing welfare device and keeping up surviving bodily function.

Also, in previous research, separation between physical and psychological load has been performed. But, our view is that there is correlation with physical and psychological load. So, we tried to measure psychological load including physical

one based on brain activity and quantitatively evaluate both load.

VI. FUTURE WORK

For the future, our purpose is to establish method of discussing useful of welfare device by evaluating load involved in other daily movements with increasing number of subjects. Especially, the investigation to relationship between load of the movement and the brain activity is important in the establishment of the evaluation method to usefulness of welfare devices and systems.

ACKNOWLEDGMENT

This research was supported as follow. There dimensional motion analysis and commodious room were provided from nac Image Technology Inc. EMG and analysis software were provided from KISSEI COMTEC. NIRS was provided from SHIMADZU COPPORATION. Developing welfare device was provided from IDEA SYSTEM CO., LTD. We are deeply grateful to them.

This study contributes to become the basis for one of theme of s-innovation program in Japan Science and Technology Agency which was named "Development Fatigue-reduction Technology for Social Contribution of Aged Person and Establishment System for Evaluation.

REFERENCES

- [1] H.Inoue, S.Shimizu, N.Tkahashi, H.Nara, T.Tsuruga, F.Miwakeichi, N.Hirai, S.Kikuchi, E. Watanabe, S.Kato," Fundamental Study to New Evaluation Method Based on Physical and Psychological Load in Care," IARIA cognitive2012, Nice, France, pp101-106.
- [2] Y. Shinoda, "Considertion of feature extraction based on center of gravity for Nihon Buyo dancer using motion capture system," SICE Annual Conference 2011, Tokyo Japan, pp. 1874-1878.
- [3] Y. Yamaguchi, A. Ishikawa, and Y.Ito, "Development of Biosignal Integration Analysis System for Human Brain Function and Behavior," Organization for Human Brain Mapping 2012, China.
- [4] H.Inoue, S. Shimizu, N. Takahashi, H.Nara, and T. Tsuruga "Fundamental Study for Evaluation of the Effect due to Exercise Load", Assistive Technology, Bio Medical Engineering and Life Support 2011, Japan.
- [5] E. Watanabe, Y. Yamashita, Y. Ito, and H. Koizumi, "Non-invasive functional mapping with multi-channel near infra-red spectroscopic topography in humans," *Heurosci Lett* 1996, Feb 16, 205(1), 41-4.
- [6] N. Takahashi, S. Shimizu, Y. Hirata, H. Nara, F. Miwakeichi, N. Hirai, S. Kikuchi, E. Watanabe, and S. Kato, "Fundamental Study for a New Assistive System during Car Driving," Proc. of International Conference on Robotics and Biomimetics, DVD-ROM, 2010, Tenjin, China.
- [7] N. Takahashi, S. Shimizu, Y. Hirata, H. Nara, H. Inoue, N. Hirai, S. Kikuchi, E. Watanabe, and S. Kato, "Basic study of Analysis of Human Brain Activities during Car Driving," the 14th International Conference on Human-Computer Interaction, 2011, USA.
- [8] S. Shimizu, N. Takahashi, H. Nara, H. Inoue, and Y. Hirata, "Fundamental Study for Human Brain Activity Based on the Spatial Cognitive Task," the 2011 Internatinal Conference on Brain Informatics-BI 2011, China.
- [9] S. Shimizu, N. Takahashi, H. Nara, H. Inoue, and Y. Hirata, "Basic Study for Human Brain Activity Based on the Spatial Cognitive Task," The Third International Conference on Advanced Cognitive Technologies and Applications, 2011, Italy.
- [10] S. Shimizu, N. Takahashi, H. Inoue, H. Nara, F. Miwakeichi, N. Hirai, S. Kikuchi, E. Watanabe, and S. Kato, "Basic Study for a New Assitive

- System Based on Brain Activity associated with Spatial Perception Task during Car driving,” Proc. International Conference on Robotics and Biomimetics, 2011, Thailand.
- [11] Y. Watanabe, “Molecular/neural mechanisms of fatigue, and the way to overcome fatigue,” *Folia Pharmacologica Japonica*, vol. 129, pp. 94-98, 2007.
- [12] H. Kuratsune, K. Yamaguti, G. Lindh, B. Evengard, G. Hagberg, K. Matsumura, M. Iwase, H. Onoe, M. Takahashi, T. Machii, Y. Kanakura, T. Kitani, B. Langstrom, and Y. Watanabe, “Brain Regions Involved in Fatigue Sensation: Reduced Acetylcarnitine Uptake in the Brain,” *Neuroimage*, vol. 17, pp. 1256-1265, November 2001.
- [13] K. Maruta, “The influence of Seat Angle on Forward Trunk Inclination During Sit-to-Stand,” *Journal of Japanese Physical Therapy Association*, vol. 31, No.1, pp. 21-28, 2004.